



Amplitude analysis and measurement of the time-dependent CP asymmetry of $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays

BABAR Collaboration ; Snoek, H L

Abstract: We present the first results on the Dalitz-plot structure and improved measurements of the time-dependent CP -violation parameters of the process $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ obtained using 468×10^6 BB decays collected with the BABAR detector at the PEP-II asymmetric-energy B factory at SLAC. The Dalitz-plot structure is probed by a time-integrated amplitude analysis that does not distinguish between B^0 and \bar{B}^0 decays. We measure the total inclusive branching fraction $B(B^0 \rightarrow K_S^0 K_S^0 K_S^0) = (6.19 \pm 0.48 \pm 0.15 \pm 0.12) \times 10^{-6}$, where the first uncertainty is statistical, the second is systematic, and the third represents the Dalitz-plot signal model dependence. We also observe evidence for the intermediate resonant states $f_0(980)$, $f_0(1710)$, and $f_2(2010)$. Their respective product branching fractions are measured to be $(2.70 - 1.19 + 1.25 \pm 0.36 \pm 1.17) \times 10^{-6}$, $(0.50 - 0.24 + 0.46 \pm 0.04 \pm 0.10) \times 10^{-6}$, and $(0.54 - 0.20 + 0.21 \pm 0.03 \pm 0.52) \times 10^{-6}$. Additionally, we determine the mixing-induced CP -violation parameters to be $S = -0.94 - 0.21 + 0.24 \pm 0.06$ and $C = -0.17 \pm 0.18 \pm 0.04$, where the first uncertainty is statistical and the second is systematic. These values are in agreement with the standard model expectation. For the first time, we report evidence of CP violation in $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays; CP conservation is excluded at 3.8 standard deviations including systematic uncertainties.

DOI: <https://doi.org/10.1103/PhysRevD.85.054023>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-75131>

Journal Article

Originally published at:

BABAR Collaboration; Snoek, H L (2012). Amplitude analysis and measurement of the time-dependent CP asymmetry of $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays. *Physical Review D (Particles, Fields, Gravitation and Cosmology)*, 85(5):054023.

DOI: <https://doi.org/10.1103/PhysRevD.85.054023>

Amplitude analysis and measurement of the time-dependent CP asymmetry of $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays

J. P. Lees, V. Poireau, and V. Tisserand

*Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),
Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France*

J. Garra Tico and E. Grauges

Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

M. Martinelli^{ab}, D. A. Milanes^a, A. Palano^{ab}, and M. Pappagallo^{ab}

INFN Sezione di Bari^a; Dipartimento di Fisica, Università di Bari^b, I-70126 Bari, Italy

G. Eigen and B. Stugu

University of Bergen, Institute of Physics, N-5007 Bergen, Norway

D. N. Brown, L. T. Kerth, Yu. G. Kolomensky, and G. Lynch

Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

H. Koch and T. Schroeder

Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

D. J. Asgeirsson, C. Hearty, T. S. Mattison, and J. A. McKenna

University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

A. Khan

Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

V. E. Blinov, A. R. Buzykaev, V. P. Druzhinin, V. B. Golubev, E. A. Kravchenko, A. P. Onuchin,

S. I. Serednyakov, Yu. I. Skovpen, E. P. Solodov, K. Yu. Todyshev, and A. N. Yushkov

Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

M. Bondioli, D. Kirkby, A. J. Lankford, M. Mandelkern, and D. P. Stoker

University of California at Irvine, Irvine, California 92697, USA

H. Atmacan, J. W. Gary, F. Liu, O. Long, and G. M. Vitug

University of California at Riverside, Riverside, California 92521, USA

C. Campagnari, T. M. Hong, D. Kovalskyi, J. D. Richman, and C. A. West

University of California at Santa Barbara, Santa Barbara, California 93106, USA

A. M. Eisner, J. Kroseberg, W. S. Lockman, A. J. Martinez, T. Schalk, B. A. Schumm, and A. Seiden

University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

C. H. Cheng, D. A. Doll, B. Echenard, K. T. Flood, D. G. Hitlin, P. Ongmongkolkul, F. C. Porter, and A. Y. Rakitin

California Institute of Technology, Pasadena, California 91125, USA

R. Andreassen, M. S. Dubrovin, Z. Huard, B. T. Meadows, M. D. Sokoloff, and L. Sun

University of Cincinnati, Cincinnati, Ohio 45221, USA

P. C. Bloom, W. T. Ford, A. Gaz, M. Nagel, U. Nauenberg, J. G. Smith, and S. R. Wagner

University of Colorado, Boulder, Colorado 80309, USA

R. Ayad* and W. H. Toki

Colorado State University, Fort Collins, Colorado 80523, USA

B. Spaan

Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany

M. J. Kobel, K. R. Schubert, and R. Schwierz

Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

D. Bernard and M. Verderi

Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France

P. J. Clark and S. Playfer

University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

D. Bettoni^a, C. Bozzi^a, R. Calabrese^{ab}, G. Cibinetto^{ab}, E. Fioravanti^{ab}, I. Garzia^{ab},

E. Luppi^{ab}, M. Menerato^{ab}, M. Negrini^{ab}, L. Piemontese^a, and V. Santoro^a

INFN Sezione di Ferrara^a; Dipartimento di Fisica, Università di Ferrara^b, I-44100 Ferrara, Italy

R. Baldini-Ferrolì, A. Calcaterra, R. de Sangro, G. Finocchiaro,

M. Nicolaci, P. Patteri, I. M. Peruzzi,[†] M. Piccolo, M. Rama, and A. Zallo

INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

R. Contri^{ab}, E. Guido^{ab}, M. Lo Vetere^{ab}, M. R. Monge^{ab}, S. Passaggio^a, C. Patrignani^{ab}, and E. Robutti^a

INFN Sezione di Genova^a; Dipartimento di Fisica, Università di Genova^b, I-16146 Genova, Italy

B. Bhuyan and V. Prasad

Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India

C. L. Lee and M. Morii

Harvard University, Cambridge, Massachusetts 02138, USA

A. J. Edwards

Harvey Mudd College, Claremont, California 91711

A. Adametz, J. Marks, and U. Uwer

Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

F. U. Bernlochner, M. Ebert, H. M. Lacker, and T. Lueck

Humboldt-Universität zu Berlin, Institut für Physik, Newtonstr. 15, D-12489 Berlin, Germany

P. D. Dauncey and M. Tibbetts

Imperial College London, London, SW7 2AZ, United Kingdom

P. K. Behera and U. Mallik

University of Iowa, Iowa City, Iowa 52242, USA

C. Chen, J. Cochran, W. T. Meyer, S. Prell, E. I. Rosenberg, and A. E. Rubin

Iowa State University, Ames, Iowa 50011-3160, USA

A. V. Gritsan and Z. J. Guo

Johns Hopkins University, Baltimore, Maryland 21218, USA

N. Arnaud, M. Davier, G. Grosdidier, F. Le Diberder, A. M. Lutz,

B. Malaescu, P. Roudeau, M. H. Schune, A. Stocchi, and G. Wormser

Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,

Centre Scientifique d'Orsay, B. P. 34, F-91898 Orsay Cedex, France

D. J. Lange and D. M. Wright

Lawrence Livermore National Laboratory, Livermore, California 94550, USA

I. Bingham, C. A. Chavez, J. P. Coleman, J. R. Fry, E. Gabathuler, D. E. Hutchcroft, D. J. Payne, and C. Touramanis
University of Liverpool, Liverpool L69 7ZE, United Kingdom

A. J. Bevan, F. Di Lodovico, R. Sacco, and M. Sigamani
Queen Mary, University of London, London, E1 4NS, United Kingdom

G. Cowan
University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

D. N. Brown and C. L. Davis
University of Louisville, Louisville, Kentucky 40292, USA

A. G. Denig, M. Fritsch, W. Gradl, A. Hafner, and E. Prencipe
Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany

K. E. Alwyn, D. Bailey, R. J. Barlow,[‡] G. Jackson, and G. D. Lafferty
University of Manchester, Manchester M13 9PL, United Kingdom

E. Behn, R. Cenci, B. Hamilton, A. Jawahery, D. A. Roberts, and G. Simi
University of Maryland, College Park, Maryland 20742, USA

C. Dallapiccola
University of Massachusetts, Amherst, Massachusetts 01003, USA

R. Cowan, D. Dujmic, and G. Sciolla
Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA

D. Lindemann, P. M. Patel, S. H. Robertson, and M. Schram
McGill University, Montréal, Québec, Canada H3A 2T8

P. Biassoni^{ab}, A. Lazzaro^{ab}, V. Lombardo^a, N. Neri^{ab}, F. Palombo^{ab}, and S. Stracka^{ab}
INFN Sezione di Milano^a; Dipartimento di Fisica, Università di Milano^b, I-20133 Milano, Italy

L. Cremaldi, R. Godang,[§] R. Kroeger, P. Sonnek, and D. J. Summers
University of Mississippi, University, Mississippi 38677, USA

X. Nguyen and P. Taras
Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7

G. De Nardo^{ab}, D. Monorchio^{ab}, G. Onorato^{ab}, and C. Sciacca^{ab}
*INFN Sezione di Napoli^a; Dipartimento di Scienze Fisiche,
 Università di Napoli Federico II^b, I-80126 Napoli, Italy*

G. Raven and H. L. Snoek
NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands

C. P. Jessop, K. J. Knoepfel, J. M. LoSecco, and W. F. Wang
University of Notre Dame, Notre Dame, Indiana 46556, USA

K. Honscheid and R. Kass
Ohio State University, Columbus, Ohio 43210, USA

J. Brau, R. Frey, N. B. Sinev, D. Strom, and E. Torrence
University of Oregon, Eugene, Oregon 97403, USA

E. Feltresi^{ab}, N. Gagliardi^{ab}, M. Margoni^{ab}, M. Morandin^a,

M. Posocco^a, M. Rotondo^a, F. Simonetto^{ab}, and R. Stroili^{ab}
INFN Sezione di Padova^a; Dipartimento di Fisica, Università di Padova^b, I-35131 Padova, Italy

S. Akar, E. Ben-Haim, M. Bomben, G. R. Bonneaud, H. Briand, G. Calderini,
 J. Chauveau, O. Hamon, Ph. Leruste, G. Marchiori, J. Ocariz, and S. Sitt
*Laboratoire de Physique Nucléaire et de Hautes Energies,
 IN2P3/CNRS, Université Pierre et Marie Curie-Paris6,
 Université Denis Diderot-Paris7, F-75252 Paris, France*

M. Biasini^{ab}, E. Manoni^{ab}, S. Pacetti^{ab}, and A. Rossi^{ab}
INFN Sezione di Perugia^a; Dipartimento di Fisica, Università di Perugia^b, I-06100 Perugia, Italy

C. Angelini^{ab}, G. Batignani^{ab}, S. Bettarini^{ab}, M. Carpinelli^{ab}, G. Casarosa^{ab}, A. Cervelli^{ab}, F. Forti^{ab},
 M. A. Giorgi^{ab}, A. Lusiani^{ac}, B. Oberhof^{ab}, E. Paoloni^{ab}, A. Perez^a, G. Rizzo^{ab}, and J. J. Walsh^a
INFN Sezione di Pisa^a; Dipartimento di Fisica, Università di Pisa^b; Scuola Normale Superiore di Pisa^c, I-56127 Pisa, Italy

D. Lopes Pegna, C. Lu, J. Olsen, A. J. S. Smith, and A. V. Telnov
Princeton University, Princeton, New Jersey 08544, USA

F. Anulli^a, G. Cavoto^a, R. Faccini^{ab}, F. Ferrarotto^a, F. Ferroni^{ab}, L. Li Gioi^a, M. A. Mazzoni^a, and G. Piredda^a
*INFN Sezione di Roma^a; Dipartimento di Fisica,
 Università di Roma La Sapienza^b, I-00185 Roma, Italy*

C. Büniger, O. Grünberg, T. Hartmann, T. Leddig, H. Schröder, and R. Waldi
Universität Rostock, D-18051 Rostock, Germany

T. Adye, E. O. Olaiya, and F. F. Wilson
Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom

S. Emery, G. Hamel de Monchenault, G. Vasseur, and Ch. Yèche
CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France

D. Aston, D. J. Bard, R. Bartoldus, C. Cartaro, M. R. Convery, J. Dorfan, G. P. Dubois-Felsmann, W. Dunwoodie,
 R. C. Field, M. Franco Sevilla, B. G. Fulsom, A. M. Gabareen, M. T. Graham, P. Grenier, C. Hast, W. R. Innes,
 M. H. Kelsey, H. Kim, P. Kim, M. L. Kocian, D. W. G. S. Leith, P. Lewis, S. Li, B. Lindquist, S. Luitz, V. Luth,
 H. L. Lynch, D. B. MacFarlane, D. R. Muller, H. Neal, S. Nelson, I. Ofte, M. Perl, T. Pulliam, B. N. Ratcliff,
 A. Roodman, A. A. Salnikov, R. H. Schindler, A. Snyder, D. Su, M. K. Sullivan, J. Va'vra, A. P. Wagner,
 M. Weaver, W. J. Wisniewski, M. Wittgen, D. H. Wright, H. W. Wulsin, A. K. Yarritu, C. C. Young, and V. Ziegler
SLAC National Accelerator Laboratory, Stanford, California 94309 USA

W. Park, M. V. Purohit, R. M. White, and J. R. Wilson
University of South Carolina, Columbia, South Carolina 29208, USA

A. Randle-Conde and S. J. Sekula
Southern Methodist University, Dallas, Texas 75275, USA

M. Bellis, J. F. Benitez, P. R. Burchat, and T. S. Miyashita
Stanford University, Stanford, California 94305-4060, USA

M. S. Alam and J. A. Ernst
State University of New York, Albany, New York 12222, USA

R. Gorodeisky, N. Guttman, D. R. Peimer, and A. Soffer
Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel

P. Lund and S. M. Spanier
University of Tennessee, Knoxville, Tennessee 37996, USA

R. Eckmann, J. L. Ritchie, A. M. Ruland, C. J. Schilling, R. F. Schwitters, and B. C. Wray
University of Texas at Austin, Austin, Texas 78712, USA

J. M. Izen and X. C. Lou
University of Texas at Dallas, Richardson, Texas 75083, USA

F. Bianchi^{ab} and D. Gamba^{ab}
INFN Sezione di Torino^a; Dipartimento di Fisica Sperimentale, Università di Torino^b, I-10125 Torino, Italy

L. Lanceri^{ab} and L. Vitale^{ab}
INFN Sezione di Trieste^a; Dipartimento di Fisica, Università di Trieste^b, I-34127 Trieste, Italy

F. Martinez-Vidal and A. Oyanguren
IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain

H. Ahmed, J. Albert, Sw. Banerjee, H. H. F. Choi, G. J. King, R. Kowalewski,
 M. J. Lewczuk, I. M. Nugent, J. M. Roney, R. J. Sobie, and N. Tasneem
University of Victoria, Victoria, British Columbia, Canada V8W 3P6

T. J. Gershon, P. F. Harrison, T. E. Latham, and E. M. T. Puccio
Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

H. R. Band, S. Dasu, Y. Pan, R. Prepost, and S. L. Wu
University of Wisconsin, Madison, Wisconsin 53706, USA

We present the first results on the Dalitz-plot structure and improved measurements of the time-dependent CP -violation parameters of the process $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ obtained using 468×10^6 $B\bar{B}$ decays collected with the BABAR detector at the PEP-II asymmetric-energy B factory at SLAC. The Dalitz-plot structure is probed by a time-integrated amplitude analysis that does not distinguish between B^0 and \bar{B}^0 decays. We measure the total inclusive branching fraction $\mathcal{B}(B^0 \rightarrow K_S^0 K_S^0 K_S^0) = (6.19 \pm 0.48 \pm 0.15 \pm 0.12) \times 10^{-6}$, where the first uncertainty is statistical, the second is systematic, and the third represents the Dalitz-plot signal model dependence. We also observe evidence for the intermediate resonant states $f_0(980)$, $f_0(1710)$, and $f_2(2010)$. Their respective product branching fractions are measured to be $(2.70_{-1.19}^{+1.25} \pm 0.36 \pm 1.17) \times 10^{-6}$, $(0.50_{-0.24}^{+0.46} \pm 0.04 \pm 0.10) \times 10^{-6}$, and $(0.54_{-0.20}^{+0.21} \pm 0.03 \pm 0.52) \times 10^{-6}$. Additionally, we determine the mixing-induced CP -violation parameters to be $\mathcal{S} = -0.94_{-0.21}^{+0.24} \pm 0.06$ and $\mathcal{C} = -0.17 \pm 0.18 \pm 0.04$, where the first uncertainty is statistical and the second is systematic. These values are in agreement with the standard model expectation. For the first time, we report evidence of CP violation in $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays; CP conservation is excluded at 3.8 standard deviations including systematic uncertainties.

PACS numbers: 13.66.Bc, 14.40.-n, 13.25.Gv, 13.25.Jx, 13.20.Jf

I. INTRODUCTION

Over the past ten years, the B factories have shown that the Cabibbo-Kobayashi-Maskawa paradigm in the standard model (SM), with a single weak phase in the quark mixing matrix, accounts for the observed CP -

symmetry violation in the quark sector. However, there may be other CP -violating sources beyond the SM. Charmless hadronic B decays, like $B^0 \rightarrow K_S^0 K_S^0 K_S^0$, are of great interest because they are dominated by loop diagrams and are thus sensitive to new physics effects at large energy scales [1]. In the SM, the mixing-induced CP -violation parameters in this decay are expected to be the same, up to $\sim 1\%$ [2], as in the tree-diagram-dominated modes such as $B^0 \rightarrow J/\psi K_S^0$. Both BABAR [3] and Belle [4] have previously performed time-dependent CP -violation measurements of the inclusive mode $B^0 \rightarrow K_S^0 K_S^0 K_S^0$, which is permissible because the final state is CP definite [5].

The structure of the Dalitz plot (DP), however, is of interest; although the time-dependent CP -violation parameters \mathcal{S} and \mathcal{C} [see Eq. (33)] can be measured inclusively without taking into account the phase space, dif-

*Now at Temple University, Philadelphia, Pennsylvania 19122, USA

†Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡Now at the University of Huddersfield, Huddersfield HD1 3DH, UK

§Now at University of South Alabama, Mobile, Alabama 36688, USA

¶Also with Università di Sassari, Sassari, Italy

ferent resonant contributions may have different values of these parameters in the presence of new physics. The statistical precision is not sufficient to perform a time-dependent amplitude analysis, but as we show below, it is possible to extract branching fractions from resonant contributions to the decay using a time-integrated amplitude analysis. Additionally, the amplitude analysis could shed light on the controversial $f_X(1500)$ resonance: recent measurements of $B^0 \rightarrow K^+K^-K_S^0$ and $B^\pm \rightarrow K^+K^-K^\pm$ from *BABAR* [6–8] and *Belle* [9, 10] have shown evidence of a wide structure in the m_{K+K^-} spectrum around 1.5 GeV. In these measurements, it was assumed that this structure is a single scalar resonance; however a vector hypothesis could not be ruled out. The *BABAR* measurement of $B^+ \rightarrow K^+K^-\pi^+$ [11] appears to show an enhancement around 1.5 GeV, while the *BABAR* analysis of $B^\pm \rightarrow K_S^0K_S^0\pi^\pm$ [12] finds no evidence of a possible $f_X(1500)$, suggesting that the structure is either a vector meson or something exotic. An amplitude analysis of $B^0 \rightarrow K_S^0K_S^0K_S^0$ will provide further insight into the nature of this structure, as only intermediate states of even spin are permitted due to Bose-Einstein statistics; an observation of the $f_X(1500)$ decaying to $K_S^0K_S^0$ would require an even-spin state. Finally, the amplitude analyses of $B \rightarrow K\pi\pi$ and $B \rightarrow KKK$ modes may be used to extract the Cabibbo-Kobayashi-Maskawa angle γ [13].

This paper presents the first amplitude analysis and the final *BABAR* update of the time-dependent CP -asymmetry measurement of $B^0 \rightarrow K_S^0K_S^0K_S^0$ using the full $\Upsilon(4S)$ dataset. The amplitude analysis is time-integrated CP -averaged (i.e., it does not use flavor-tagging information to distinguish between B^0 and \bar{B}^0 mesons). It takes advantage of the interference pattern in the DP to measure relative magnitudes and phases for the different resonant modes using $B^0 \rightarrow K_S^0K_S^0K_S^0$ decays with $K_S^0 \rightarrow \pi^+\pi^-$, denoted by $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$. The magnitudes and phases are then translated into individual branching fractions for the resonant modes. The time-dependent analysis extracts the S and C parameters by modeling the proper-time distribution. This part of the analysis uses both $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ events and events where one of the K_S^0 mesons decays to $\pi^0\pi^0$, denoted by $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$.

In Sec. II we briefly describe the *BABAR* detector and the dataset. The amplitude analysis is described in Sec. III and the time-dependent analysis in Sec. IV. Finally we summarize the results in Sec. V.

II. THE *BABAR* DETECTOR AND DATASET

The data used in this analysis were collected with the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- storage ring at SLAC. The sample consists of an integrated luminosity of 426.0 fb^{-1} , corresponding to $(467.8 \pm 5.1) \times 10^6 B\bar{B}$ pairs collected at the $\Upsilon(4S)$ resonance (“on-resonance”), and 44.5 fb^{-1} collected about

40 MeV below the $\Upsilon(4S)$ (“off-resonance”).

A detailed description of the *BABAR* detector is presented in Ref. [14]. The tracking system used for track and vertex reconstruction has two components: a silicon vertex tracker and a drift chamber, both operating within a 1.5 T magnetic field generated by a superconducting solenoidal magnet. A detector of internally reflected Cherenkov light associates Cherenkov photons with tracks for particle identification. The energies of photons and electrons are determined from the measured light produced in electromagnetic showers inside a CsI crystal electromagnetic calorimeter. Muon candidates are identified with the use of the instrumented flux return of the solenoid.

III. AMPLITUDE ANALYSIS

In Secs. III A and III B we describe the DP formalism and introduce the signal parameters that are extracted from data. In Sec. III C we describe the requirements used to select the signal candidates and suppress backgrounds. In Sec. III D we describe the fit method and the approach used to account for experimental effects such as resolution. In Sec. III E we present the results of the fit, and finally, in Sec. III F we discuss systematic uncertainties in the results.

A. Decay amplitudes

The $B^0 \rightarrow K_S^0K_S^0K_S^0$ decay contains three identical particles in the final state and therefore the amplitude needs to be symmetrized. We consider the decay of a spin-zero B^0 into three daughters, $K_S^0(1)$, $K_S^0(2)$, and $K_S^0(3)$, with four-momenta p_1 , p_2 , and p_3 . The decay amplitude is given by [2]

$$\begin{aligned} \mathcal{A}[B^0 \rightarrow K_S^0(1)K_S^0(2)K_S^0(3)] & \\ &= \left(\frac{1}{2}\right)^{3/2} \left\{ \mathcal{A}_1[B^0 \rightarrow \bar{K}^0(1)K^0(2)K^0(3)] \right. \\ &\quad + \mathcal{A}_2[B^0 \rightarrow \bar{K}^0(2)K^0(3)K^0(1)] \\ &\quad \left. + \mathcal{A}_3[B^0 \rightarrow \bar{K}^0(3)K^0(1)K^0(2)] \right\}, \end{aligned} \quad (1)$$

which takes into account the three permitted paths from the initial state to the final state. For instance for the B^0 decay this consists of an intermediate state $K^0K^0\bar{K}^0$. Since the labeling of the three identical particles is arbitrary, we classify the final-state particles according to the square of the invariant mass, s_{ij} , defined as

$$s_{ij} = s_{ji} = m_{K_S^0(i)K_S^0(j)}^2 = (p_i + p_j)^2, \quad (2)$$

where i and j are the K_S^0 indices. We use as independent (Mandelstam) variables the minimum and the maximum

of the squared masses s_{\min} and s_{\max} :

$$\begin{aligned} s_{\min} &= \min(s_{12}, s_{23}, s_{13}), \\ s_{\max} &= \max(s_{12}, s_{23}, s_{13}). \end{aligned} \quad (3)$$

The third (median) invariant squared mass s_{med} can be obtained from energy and momentum conservation:

$$s_{\text{med}} = m_{B^0}^2 + 3m_{K_S^0}^2 - s_{\min} - s_{\max}. \quad (4)$$

The differential B meson decay width with respect to the variables defined in Eq. (3) (i.e., the DP variables) reads

$$d\Gamma(B \rightarrow K_S^0 K_S^0 K_S^0) = \frac{1}{(2\pi)^3} \frac{|\mathcal{A}|^2}{32m_{B^0}^3} ds_{\min} ds_{\max}, \quad (5)$$

where \mathcal{A} is the Lorentz-invariant amplitude of the three-body decay. This amplitude analysis does not take into account any flavor tagging or time dependence, thus it is CP averaged and time integrated. The term $|\mathcal{A}|^2$ is therefore simply the average of squares of the contributions $\mathcal{A}[B^0 \rightarrow K_S^0 K_S^0 K_S^0]$ and $\mathcal{A}[\bar{B}^0 \rightarrow K_S^0 K_S^0 K_S^0]$.

The choice of the variables s_{\min} and s_{\max} gives a uniquely defined coordinate in the symmetrized DP. Therefore only one sixth of the DP is populated, i.e., the event density is six times larger compared to an amplitude analysis involving three distinct particles.

We describe the distribution of signal events in the DP using an isobar approximation, which models the total amplitude as resulting from a coherent sum of amplitudes from the N individual decay channels of the B meson, either into an intermediate resonance and a bachelor particle or in a nonresonant manner:

$$\mathcal{A}(s_{\min}, s_{\max}) = \sum_{j=1}^N c_j F_j(s_{\min}, s_{\max}). \quad (6)$$

Here F_j (described in detail below) are DP-dependent amplitudes containing the decay dynamics and c_j are complex coefficients describing the relative magnitudes and phases of the different decay channels. This description, which contains a single complex number c_j for each decay channel regardless of the B flavor (B^0 or \bar{B}^0), reflects the assumptions of no direct CP violation and of a common weak phase for all the decay channels. With this description we cannot extract any weak phase information; this would require using per- B flavor complex amplitudes. The factor F_j contains strong dynamics only, and thus does not change under CP conjugation.

Intermediate resonances decay to $K^0 \bar{K}^0$. In terms of the isobar approximation, the amplitude in Eq. (1) for a resonant state j becomes

$$\begin{aligned} \mathcal{A}[B^0 \rightarrow K_S^0(1)K_S^0(2)K_S^0(3)] \\ \propto c_j [F_j(s_{12}, s_{13}) + F_j(s_{12}, s_{23}) + F_j(s_{13}, s_{23})]. \end{aligned} \quad (7)$$

This reflects the fact that it is impossible to associate a given K_S^0 to a flavor eigenstate K^0 or \bar{K}^0 . In practice,

this sum of three F_j terms, corresponding to an even-spin resonance, is implicitly taken into account by the description in terms of s_{\min} and s_{\max} .

The F_j terms are represented by the product of the invariant mass and angular distributions; i.e.,

$$F_j(s_{\min}, s_{\max}, L) = R_j(m) X_L(|\vec{p}^*| r') X_L(|\vec{q}| r) T_j(L, \vec{p}, \vec{q}), \quad (8)$$

where

- (i) m is the invariant mass of the decay products of the resonance;
- (ii) $R_j(m)$ is the resonance mass term or “line shape”, e.g., relativistic Breit–Wigner (RBW);
- (iii) L is the orbital angular momentum between the resonance and the bachelor particle;
- (iv) \vec{p}^* is the momentum of the bachelor particle evaluated in the rest frame of the B ;
- (v) \vec{p} and \vec{q} are the momenta of the bachelor particle and one of the resonance daughters, respectively, both evaluated in the rest frame of the resonance;
- (vi) $X_L(|\vec{p}^*| r')$ and $X_L(|\vec{q}| r)$ are Blatt–Weisskopf barrier factors [15] with barrier radii of r and r' , and
- (vii) $T_j(L, \vec{p}, \vec{q})$ is the angular distribution:

$$L = 0 : T_j = 1, \quad (9)$$

$$L = 2 : T_j = \frac{8}{3} [3(\vec{p} \cdot \vec{q})^2 - (|\vec{p}| |\vec{q}|)^2]. \quad (10)$$

The Blatt–Weisskopf barrier factor is unity for all the zero-spin resonances. In our analysis it is relevant only for the $f_2(2010)$. Since for this resonance r and r' are not measured, we take them both to be 1.5 GeV^{-1} and vary by $\pm 0.5 \text{ GeV}^{-1}$ to estimate the systematic uncertainty.

The helicity angle of a resonance is defined as the angle between \vec{p} and \vec{q} . Explicitly, the helicity angle θ for a given resonance is defined between the momenta of the bachelor particle and one of the daughters of the resonance in the resonance rest frame. Because of the identical final-state particles this definition is ambiguous, but the ambiguity disappears because of the description of the DP in terms of s_{\min} and s_{\max} . There are three possible invariant-mass combinations: s_{\min} , s_{med} , and s_{\max} . We denote the corresponding helicity angles as θ_{\min} , θ_{med} , and θ_{\max} . The three angles are defined between 0 and $\pi/2$.

As the present study is the first amplitude analysis of this decay, we use the method outlined in Sec. III D 3 to determine the contributing intermediate states. The components of the nominal signal model are summarized in Table I.

For most resonances in this analysis the R_j are taken to be RBW [16] line shapes:

$$R_j(m) = \frac{1}{(m_0^2 - m^2) - im_0 \Gamma(m)}, \quad (11)$$

where m_0 is the nominal mass of the resonance and $\Gamma(m)$ is the mass-dependent width. In the general case of a spin- J resonance, the latter can be expressed as

$$\Gamma(m) = \Gamma_0 \left(\frac{q}{q_0} \right)^{2J+1} \left(\frac{m_0}{m} \right) \frac{X_J^2(|\vec{q}|r)}{X_J^2(|\vec{q}_0|r)}. \quad (12)$$

The symbol Γ_0 denotes the nominal width of the resonance. The values of m_0 and Γ_0 are listed in Table I. The symbol q_0 denotes the value of q when $m = m_0$.

For the $f_0(980)$ line shape the Flatté form [17] is used. In this case the mass-dependent width is given by the sum of the widths in the $\pi\pi$ and KK systems:

$$\Gamma(m) = \Gamma_{\pi\pi}(m) + \Gamma_{KK}(m), \quad (13)$$

where

$$\Gamma_{\pi\pi}(m) = g_\pi \left(\frac{1}{3} \sqrt{1 - 4m_{\pi^0}^2/m^2} + \frac{2}{3} \sqrt{1 - 4m_{\pi^\pm}^2/m^2} \right), \quad (14)$$

$$\Gamma_{KK}(m) = g_K \left(\frac{1}{2} \sqrt{1 - 4m_{K^\pm}^2/m^2} + \frac{1}{2} \sqrt{1 - 4m_{K^0}^2/m^2} \right). \quad (15)$$

The fractional coefficients arise from isospin conservation and g_π and g_K are coupling constants for which the values are given in Table I. The nonresonant (NR) component is modeled using an exponential function:

$$R_{\text{NR}}(m) = e^{\alpha m^2}. \quad (16)$$

As in the resonant case, here m is the invariant mass of the relevant $K_S^0 K_S^0$ pair. The parameter α is taken from the *BABAR* $B^+ \rightarrow K^+ K^- K^+$ analysis [7, 8] and is given in Table I. This value was found to be compatible with the one resulting from varying α in the maximum-likelihood fit in the present analysis. There is no satisfactory theoretical description of the NR component; it has to be determined empirically. The exponential function of Eq. (16) was used by other amplitude analyses of B -meson decays to three kaons [6–10]. Adopting the same parametrization for the NR term allows the comparison of results for other components.

B. The square Dalitz plot

We use two-dimensional histograms to describe the phase-space dependent reconstruction efficiency and to model the background over the DP. When the phase-space boundaries of the DP do not coincide with the histogram bin boundaries this may introduce biases. We

therefore define h_{\min} and h_{\max} as $\cos\theta_{\min}$ and $\cos\theta_{\max}$, respectively, and apply the transformation

$$(s_{\min}, s_{\max}) \longrightarrow (h_{\min}, h_{\max}). \quad (17)$$

The (h_{\min}, h_{\max}) plane is referred to as the square Dalitz plot (SDP), where both h_{\min} and h_{\max} range between 0 and 1 due to the convention adopted for the helicity angles (see Fig. 1). Explicitly, the transformation is

$$h_{\min} = \frac{s_{\min}(s_{\max} - s_{\text{med}})}{\sqrt{s_{\min}^2 - 4m_{K_S^0}^2 s_{\min}}} \times \frac{1}{\sqrt{(m_{B^0}^2 - m_{K_S^0}^2 - s_{\min})^2 - 4m_{K_S^0}^2 s_{\min}}}, \quad (18)$$

$$h_{\max} = \frac{s_{\max}(s_{\text{med}} - s_{\min})}{\sqrt{s_{\max}^2 - 4m_{K_S^0}^2 s_{\max}}} \times \frac{1}{\sqrt{(m_{B^0}^2 - m_{K_S^0}^2 - s_{\max})^2 - 4m_{K_S^0}^2 s_{\max}}}, \quad (19)$$

where the numerators may easily be expressed in terms of s_{\min} and s_{\max} using Eq. (4). The differential surface elements of the DP and the SDP are related by

$$ds_{\min} ds_{\max} = |\det J| dh_{\min} dh_{\max}, \quad (20)$$

where $J = J(h_{\min}, h_{\max})$ is the appropriate Jacobian matrix. The backward transformations $s_{\min}(h_{\min}, h_{\max})$ and $s_{\max}(h_{\min}, h_{\max})$, and therefore the Jacobian $|\det J|$, cannot be found analytically; they are obtained numerically. The variables h_{\min} and h_{\max} as a function of the invariant masses are shown in Fig. 1 together with the Jacobian.

C. Event selection and backgrounds

We reconstruct $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ candidates from three $K_S^0 \rightarrow \pi^+ \pi^-$ candidates that form a good quality vertex; i.e., the fit of the B^0 vertex is required to converge and the χ^2 probability of each K_S^0 vertex fit has to be greater than 10^{-6} . Each K_S^0 candidate must have $\pi^+ \pi^-$ invariant mass within 12.1 MeV/ c^2 of the nominal K^0 mass [16], and decay length with respect to the B vertex between 0.22 and 45 cm. The last criterion ensures that the decay vertices of the B^0 and the K_S^0 are well separated. In addition, combinatorial background is suppressed by selecting events for which the angle between the momentum vector of each K_S^0 candidate and the vector connecting the beamspot and the K_S^0 vertex is smaller than 0.0185 radians. We ensure a good B vertex fit quality by requiring that the charged pions of at least one of the K_S^0 candidates have hits in the two inner layers of the vertex tracker.

A B meson candidate is characterized kinematically by the energy-substituted mass $m_{\text{ES}} \equiv \sqrt{(s/2 + \vec{p}_i \cdot \vec{p}_B)^2 / E_i^2 - \vec{p}_B^2}$ and the energy difference

TABLE I: Parameters of the DP model used in the fit. The Blatt–Weisskopf barrier parameters (r and r') of the $f_2(2010)$, which have not been measured, are varied by $\pm 0.5 \text{ GeV}^{-1}$ for the model uncertainty.

Resonance	Parameters	Line shape	Reference
$f_0(980)$	$m_0 = (965 \pm 10) \text{ MeV}/c^2$	Flatté	[18]
	$g_\pi = (165 \pm 18) \text{ MeV}/c^2$	Eq. (13)	
	$g_K = (695 \pm 93) \text{ MeV}/c^2$		
$f_0(1710)$	$m_0 = (1724 \pm 7) \text{ MeV}/c^2$	RBW	[16]
	$\Gamma_0 = (137 \pm 8) \text{ MeV}/c^2$	Eq. (11)	
$f_2(2010)$	$m_0 = (2011^{+60}_{-80}) \text{ MeV}/c^2$	RBW	[16]
	$\Gamma_0 = (202 \pm 60) \text{ MeV}/c^2$	Eq. (11)	
	$r = r' = 1.5 \text{ GeV}^{-1}$		
NR decays	$\alpha = (-0.14 \pm 0.02) \text{ GeV}^{-2} c^4$	Exponential NR Eq. (16)	[8]
χ_{c0}	$m_0 = (3414.75 \pm 0.31) \text{ MeV}/c^2$	RBW	[16]
	$\Gamma_0 = (10.2 \pm 0.7) \text{ MeV}/c^2$	Eq. (11)	




FIG. 1: (color online). Lines of constant helicity angle in the Dalitz plot of s_{\min} versus s_{\max} (left), and the magnitude of the Jacobian (gray scale on the right) mapping (s_{\min}, s_{\max}) to (h_{\min}, h_{\max}) . For the latter see Eq. (20).

$\Delta E \equiv E_B^* - \frac{1}{2}\sqrt{s}$, where (E_B, \vec{p}_B) and (E_i, \vec{p}_i) are the four-vectors in the laboratory frame of the B -candidate and the initial electron-positron system, respectively, and p_B is the magnitude of \vec{p}_B . The asterisk denotes the $\Upsilon(4S)$ frame, and s is the square of the invariant mass of the electron-positron system. We require $5.27 < m_{\text{ES}} < 5.29 \text{ GeV}/c^2$ and $|\Delta E| < 0.1 \text{ GeV}$. Following the calculation of these kinematic variables, each of the B candidates is refitted with its mass constrained to the world average value of the B meson mass [16] in order to improve the DP position resolution, and ensure that Eq. (4) holds. The sideband used for background studies is in the range $5.20 < m_{\text{ES}} < 5.27 \text{ GeV}/c^2$ and $|\Delta E| < 0.1 \text{ GeV}$.

Backgrounds arise primarily from random combinations in continuum $e^+e^- \rightarrow q\bar{q}$ events ($q = d, u, s, c$). To enhance discrimination between signal and contin-

uum background, we use a neural network (NN) [19] to combine four discriminating variables: the angles with respect to the beam axis of the B momentum and B thrust axis in the $\Upsilon(4S)$ frame, and the zeroth- and second-order monomials $L_{0,2}$ of the energy flow about the B thrust axis. The monomials are defined by $L_n = \sum_i p_i \times |\cos \theta_i|^n$, where θ_i is the angle with respect to the B thrust axis of track or neutral cluster i and p_i is the magnitude of its momentum. The sum excludes the B candidate and all quantities are calculated in the $\Upsilon(4S)$ frame. The NN is trained with off-resonance data, sideband data and simulated signal events that pass the selection criteria. Approximately 0.5% of events passing the full selection have more than one candidate. When this occurs, we select the candidate for which the error-weighted average of the masses of the K_S^0 candidates is closest to the world average K_S^0 mass [16]. With the

above selection criteria, we obtain a signal reconstruction efficiency of 6.6% that has been determined from a signal Monte Carlo (MC) sample generated using the same DP model and parameters as obtained from the data fit results. We estimate from this MC that 1.4% of the selected signal events are misreconstructed, and assign a systematic uncertainty (see Sec. III F). We use MC events to study the background from other B decays (B background). We expect fewer than 6 such events in our data sample. As these events are wrongly reconstructed, the m_{ES} and ΔE distributions are continuumlike and as a result the events are mostly absorbed in the continuum background category. We assign a systematic uncertainty for B background contamination in the signal.

D. The maximum-likelihood fit

We perform an unbinned extended maximum-likelihood fit to extract the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ event yield, as well as the resonant and nonresonant amplitudes. The fit for the amplitude analysis uses the variables m_{ES} and ΔE , the NN output, and the SDP variables to discriminate signal from background. The selected on-resonance data sample is assumed to consist of signal and continuum background. The feed-through from B decays other than the signal is found to be negligible. Misreconstructed signal events are not considered as a separate event species, but are taken into account as a part of the signal. The likelihood function \mathcal{L}_i for event i is the sum

$$\mathcal{L}_i = \sum_j N_j \mathcal{P}_j^i(m_{\text{ES}}, \Delta E, \text{NN}, h_{\min}, h_{\max}), \quad (21)$$

where j stands for the species (signal, continuum background) and N_j is the corresponding yield. Each probability density function (PDF) \mathcal{P}_j^i is the product of four individual PDFs:

$$\mathcal{P}_j^i = \mathcal{P}_j^i(m_{\text{ES}}) \mathcal{P}_j^i(\Delta E) \mathcal{P}_j^i(\text{NN}) \mathcal{P}_j^i(h_{\min}, h_{\max}). \quad (22)$$

A study with fully reconstructed MC samples shows that correlations between the PDF variables are small and therefore we neglect them. However, possible small discrepancies in the fit results due to these correlations are accounted for in the systematic uncertainty (see Sec. III F). The total likelihood is given by

$$\mathcal{L} = \exp(-\sum_j N_j) \prod_i \mathcal{L}_i. \quad (23)$$

1. The m_{ES} , ΔE , and NN PDFs

The m_{ES} and ΔE distributions of signal events are parametrized by an asymmetric Gaussian with power-law tails:

$$\begin{aligned} &\text{Cr}(x; m_0, \sigma_l, \sigma_r, \alpha_l, \alpha_r) = \\ &\exp\left(-\frac{(x - m_0)^2}{2\sigma_i^2 + \alpha_i(x - m_0)^2}\right) \begin{cases} x - m_0 < 0 : i = l \\ x - m_0 \geq 0 : i = r \end{cases}. \end{aligned} \quad (24)$$

The m_0 parameters for both m_{ES} and ΔE are free in the fit to data, while the other parameters are fixed to values determined from a fit to MC simulation. For the NN distributions of signal we use a histogram PDF from MC simulation.

For continuum events the m_{ES} and ΔE PDFs are parametrized by an ARGUS shape function [20] and a straight line, respectively. The NN PDF is described by a sum of power functions:

$$\begin{aligned} E(x; c_1, a, b_0, b_1, b_2, b_3, c_2, c_3) = & \\ &\cos^2(c_1) [\cos^2(a) \mathcal{N}(b_0, b_1) x^{b_0} (1-x)^{b_1} \\ &+ \sin^2(a) \mathcal{N}(b_2, b_3) x^{b_2} (1-x)^{b_3}] \\ &+ \sin^2(c_1) \mathcal{N}(c_2, c_3) x^{c_2} (1-x)^{c_3}, \end{aligned} \quad (25)$$

where $x = (\text{NN} - \text{NN}_{\min}) / (\text{NN}_{\max} - \text{NN}_{\min})$ and the \mathcal{N} are normalization factors, computed analytically using the standard Γ function:

$$\mathcal{N}(\alpha, \beta) = \frac{\Gamma(\beta + 2 + \alpha)}{\Gamma(\alpha + 1)\Gamma(\beta + 1)}. \quad (26)$$

The parameters for all the continuum PDFs are determined by a fit to sideband data and then fixed for the fit in the signal region.

2. Dalitz-plot PDFs

The SDP PDF for continuum background is a histogram obtained from m_{ES} sideband on-resonance events. The SDP signal PDFs require as input the DP-dependent selection efficiency, $\varepsilon = \varepsilon(h_{\min}, h_{\max})$, that is described by a histogram and is taken from MC simulation. For each event we define the SDP signal PDF:

$$\mathcal{P}_{\text{sig}}^i(h_{\min}, h_{\max}) \propto \varepsilon(h_{\min}, h_{\max}) |\mathcal{A}(h_{\min}, h_{\max})|^2. \quad (27)$$

The normalization of the PDF is implemented by numerical integration. To describe the experimental resolution in the SDP variables, we use an ensemble of two-dimensional histograms that represents the probability to reconstruct at the coordinate (h_{\min}', h_{\max}') an event that has the true coordinate (h_{\min}, h_{\max}) . These histograms are taken from MC simulation and are convolved with the signal PDF.

3. Determination of the signal Dalitz-plot model

Using on-resonance data, we determine a nominal signal DP model by making likelihood scans with various combinations of isobars. We start from a baseline model that includes $f_0(980)$, χ_{c0} , and NR components. We then add another scalar resonance described by the RBW parametrization. We scan the likelihood by fixing the width and mass of this additional resonance at several consecutive values, for each of which the fit to the data

is repeated. All isobar magnitudes and phases are floating in these fits. From the scans we observe a significant improvement of the fit around a width and mass that are compatible with the values of the $f_0(1710)$ resonance [16]. After adding the $f_0(1710)$ to the nominal model we repeat the same procedure for an additional tensor particle. We find that the $f_2(2010)$ has a significant contribution. The results of the likelihood scans are shown in Fig. 2 in terms of $-2\Delta\ln\mathcal{L} = -2\ln\mathcal{L} - (-2\ln\mathcal{L})_{\min}$, where $(-2\ln\mathcal{L})_{\min}$ corresponds to the minimal value obtained in the particular scan. To conclude the search for possible resonant contributions we add all well established resonances [16] and check if the likelihood increases. We do not find any other significant resonant contribution but as we cannot exclude small contributions from the $f_0(1370)$, $f_2(1270)$, $f_2'(1525)$, $a_0(1450)$, and $f_0(1500)$ resonances, we assign model uncertainties (see Sec. III F) due to not taking these resonances into account.

E. Results

The maximum-likelihood fit of 505 candidates results in a $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ event yield of 200 ± 15 and a continuum yield of 305 ± 18 , where the uncertainties are statistical only. The symmetrized and square Dalitz plots of a signal DP-model MC sample generated with the result of the fit to data are shown in Fig. 3. Figure 4 shows plots of ΔE , m_{ES} , and the NN for isolated signal and continuum background events obtained by the *sPlots* [21] technique. Figure 5 shows projections of the data onto the invariant masses s_{\min} and s_{\max} .

When the fit is repeated with initial parameter values randomly chosen within wide ranges above and below the nominal values for the magnitudes and within the $[-\pi, \pi]$ interval for the phases, we observe convergence towards two solutions with minimum values of the negative log likelihood function $-2\ln\mathcal{L}$ separated by 3.25 units. In the following, we refer to them as Solution 1 (the global minimum) and Solution 2 (a local minimum). No other local minima were found.

In the fit, we measure directly the relative magnitudes and phases of the different components of the signal model. The magnitude and phase of the NR amplitude are fixed to 1 and 0, respectively, as a reference. In Fig. 6 we show likelihood scans of the isobar magnitudes and phases of all the resonances, where both solutions can be noticed. Each of these scans is obtained by fixing the corresponding isobar parameter at several consecutive values, for each of which the fit to the data is repeated. The measured relative amplitudes c_μ are used to extract the fit fraction defined as

$$FF(k) = \frac{\sum_{\mu=3k-2}^{3k} \sum_{\nu=3k-2}^{3k} c_\mu c_\nu^* \langle F_\mu F_\nu^* \rangle}{\sum_{\mu\nu} c_\mu c_\nu^* \langle F_\mu F_\nu^* \rangle}, \quad (28)$$

where k , which varies from 1 to 5, represents an intermediate state. Each fit fraction is a sum of three identical

contributions, one for each pair of K_S^0 . The indices μ and ν run from 1 to 15, as each of the five resonances contributes to three pairs of K_S^0 , which correspond to the three terms $(3k-2, 3k-1, \text{ and } 3k)$ in each sum in the numerator of Eq. (28). The dynamical amplitudes F are defined in Sec. III A and the terms

$$\langle F_\mu F_\nu^* \rangle = \int \int F_\mu F_\nu^* ds_{\min} ds_{\max} \quad (29)$$

are obtained by integration over the DP. The total fit fraction is defined as the algebraic sum of all fit fractions. This quantity is not necessarily unity due to the potential presence of net constructive or destructive interference.

In order to estimate the statistical significance of each resonance, we evaluate the difference $\Delta\ln\mathcal{L}$ between the log-likelihood of the nominal fit and that of a fit where the magnitude of the amplitude of the resonance is set to 0 (this difference can be directly read from the likelihood scans as a function of magnitudes in Fig. 6). In this case the phase of the resonance becomes meaningless, and we therefore account for two degrees of freedom removed from the fit. The value $2\Delta\ln\mathcal{L}$ is used to evaluate the p -value for 2 degrees of freedom; we determine the equivalent one-dimensional significance from this p -value.

The results for the phase and the fit fraction are given in Table II for the two solutions; the change in likelihood when the amplitude of the resonance is set to 0 and the resulting statistical significance of each resonance is given for Solution 1.

As the fit fractions are not parameters of the PDF itself, their statistical errors are obtained from the 68.3% coverage intervals of the fit-fraction distributions obtained from a large number of pseudoexperiments generated with the corresponding solution (1 or 2). As observed in other three-kaon modes [6–10], the total FF significantly exceeds unity.

In Table II it can be seen that the two solutions differ mostly in the fraction assigned to the NR and the $f_0(980)$ components. Solution 1 corresponds to a small FF of the $f_0(980)$ and a large value for the NR, and Solution 2 has a large $f_0(980)$ fraction and a smaller NR fraction. Other three-kaon modes [6–10] favor the behavior of Solution 1.

Generalizing Eq. (28), we obtain the interference fractions among the intermediate decay modes k and j

$$FF(k, j) = \frac{\sum_{\mu=3k-2}^{3k} \sum_{\nu=3j-2}^{3j} c_\mu c_\nu^* \langle F_\mu F_\nu^* \rangle}{\sum_{\mu\nu} c_\mu c_\nu^* \langle F_\mu F_\nu^* \rangle}, \quad (30)$$

which are given in Table III for Solution 1. Unlike the total FF defined above, the elements of this matrix sum to unity. The large destructive interference between the $f_0(980)K_S^0$ and the NR components appears clearly in the table. This is possible due to the large overlap in phase space between the exponential NR term and the broad tail of the $f_0(980)$ resonance above the KK threshold.

Using the relative fit fractions, we calculate the branching fraction \mathcal{B} for the intermediate mode k as

$$FF(k) \times \mathcal{B}(B^0 \rightarrow K_S^0 K_S^0 K_S^0), \quad (31)$$




FIG. 2: Two-dimensional scans of $-2\Delta\ln\mathcal{L}$ (gray scale) as a function of the mass and the width of an additional resonance. These scans were performed to look for an additional scalar resonance (left) and an additional tensor resonance (right). The baseline model of the scans for additional scalar resonances contains $f_0(980)$, χ_{c0} , and NR intermediate states. The baseline model of the scans for additional tensor resonances contains $f_0(980)$, χ_{c0} , NR, and $f_0(1710)$ intermediate states. The ellipses indicate the world average parameters [16] for the $f_0(1710)$ and $f_2(2010)$ resonances that are added to the model.




FIG. 3: Symmetrized (left) and square (right) DP for MC simulated signal events using the amplitudes obtained from the fit to data. The low population in bins along the edge of the symmetrized DP is due to the fact that the phase space boundaries do not coincide with the histogram bin boundaries.

where $\mathcal{B}(B^0 \rightarrow K_S^0 K_S^0 K_S^0)$ is the total inclusive branching fraction:

$$\mathcal{B}(B^0 \rightarrow K_S^0 K_S^0 K_S^0) = \frac{N_{\text{sig}}}{\bar{\epsilon} N_{B\bar{B}}}. \quad (32)$$

We estimate the average efficiency $\bar{\epsilon} = 6.6\%$ using a fully reconstructed DP-model MC sample generated with the parameters found in data. The results of the branching fraction measurements are shown in Table IV. As a cross check we attempt to compare our measured branching fractions to results from other measurements; however, many of the branching fractions for the decay into kaons of the resonances included in our model are not (or are only poorly) measured (marked as “seen” in Ref. [16]).

An exception is the charmonium state χ_{c0} , for which the measured value is $\mathcal{B}(\chi_{c0} \rightarrow K_S^0 K_S^0) = (3.16 \pm 0.18) \times 10^{-3}$ [16]. We can then use the *BABAR* measurement of $\mathcal{B}(B^0 \rightarrow \chi_{c0} K^0) = (142^{+55}_{-44} \pm 8 \pm 16 \pm 12) \times 10^{-6}$ [22] to calculate $\mathcal{B}[B^0 \rightarrow \chi_{c0}(\rightarrow K_S^0 K_S^0) K_S^0] = \frac{1}{2} \mathcal{B}(B^0 \rightarrow \chi_{c0} K^0) \times \mathcal{B}(\chi_{c0} \rightarrow K_S^0 K_S^0) = (0.224 \pm 0.078) \times 10^{-6}$, which is consistent with our measured branching fraction, given in Table IV.

An interesting conclusion from this first amplitude analysis of the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decay mode is that we do not need to include a broad scalar $f_X(1500)$ resonance, as has been done in other measurements [6–10], to describe the data. The peak in the invariant mass between 1.5




FIG. 4: (color online). $sPlots$ (points with error bars) and PDFs (histograms) of the discriminating variables: m_{ES} (top), ΔE (middle), and NN (bottom), for signal events (left) and continuum events (right). Below each bin are shown the residuals, normalized in error units. The horizontal dotted and full lines mark the one- and two-standard deviation levels, respectively.

and $1.6 \text{ GeV}/c^2$ can be described by the interference between the $f_0(1710)$ resonance and the nonresonant component. However minor contributions from the $f'_2(1525)$ and $f_0(1500)$ resonances to this structure cannot be excluded.

F. Systematic uncertainties

Systematic effects are divided into model and experimental uncertainties. Details on how they have been estimated are given below and the associated numerical values are summarized in Table V.

1. Model uncertainties

We vary the mass, width, and any other parameter of all isobar fit components within their errors, as quoted in Table I, and assign the observed differences in our observables as the first part of the model uncertainty (“Model” in Table V). To estimate the contribution to $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ from resonances that are not included in our signal model but cannot be excluded statistically, namely the $f_0(1370)$, $f_2(1270)$, $f'_2(1525)$, $a_0(1450)$, and $f_0(1500)$ resonances, we perform fits to pseudoexperiments that include these resonances. The masses and the widths are taken from [16], except for the $f_0(1370)$ for which we take the values from [23]. We generate pseu-




FIG. 5: (color online). Projections onto $\sqrt{s_{\min}}$ (left) and $\sqrt{s_{\max}}$ (right). On-resonance data are shown as points with error bars while the dashed (dotted) histogram represents the signal (continuum) component. The solid-line histogram is the total PDF. Below each bin are shown the residuals, normalized in error units. The horizontal dotted and full lines mark the one- and two-standard deviation levels, respectively.

TABLE II: Summary of measurements of the quasi-two-body parameters. The quoted uncertainties are statistical only. The change in the log-likelihood ($-2\Delta \ln \mathcal{L}$) corresponds to the case where the magnitude of the amplitude of the resonance is set to 0. This number is used for the estimation of the statistical significance of each resonance.

Mode	Parameter	Solution 1	Solution 2
$f_0(980)K_S^0$	FF	$0.44^{+0.20}_{-0.19}$	$1.03^{+0.22}_{-0.17}$
	Phase [rad]	0.09 ± 0.16	1.26 ± 0.17
	$-2\Delta \ln \mathcal{L}$	11.7	-
	Significance [σ]	3.0	-
$f_0(1710)K_S^0$	FF	$0.07^{+0.07}_{-0.03}$	$0.09^{+0.05}_{-0.02}$
	Phase [rad]	1.11 ± 0.23	0.36 ± 0.20
	$-2\Delta \ln \mathcal{L}$	14.2	-
	Significance [σ]	3.3	-
$f_2(2010)K_S^0$	FF	$0.09^{+0.03}_{-0.03}$	0.10 ± 0.02
	Phase [rad]	2.50 ± 0.20	1.58 ± 0.22
	$-2\Delta \ln \mathcal{L}$	14.0	-
	Significance [σ]	3.3	-
NR	FF	$2.16^{+0.36}_{-0.37}$	$1.37^{+0.26}_{-0.21}$
	Phase [rad]	0.0	0.0
	$-2\Delta \ln \mathcal{L}$	68.1	-
	Significance [σ]	8.0	-
$\chi_{c0}K_S^0$	FF	$0.07^{+0.04}_{-0.02}$	0.07 ± 0.02
	Phase [rad]	0.63 ± 0.47	-0.24 ± 0.52
	$-2\Delta \ln \mathcal{L}$	18.5	-
	Significance [σ]	3.9	-
Total FF		$2.84^{+0.71}_{-0.66}$	$2.66^{+0.35}_{-0.27}$

doexperiments with the additional resonances, where the isobar magnitudes and phases have been determined in fits to data, and fit these datasets with the nominal model. We assign the induced shift in the observables as a second part of the model uncertainty.

2. Experimental systematic uncertainties

To validate the analysis procedure, we perform fits on a large number of pseudoexperiments generated with the measured yields of signal events and continuum background. The signal events are taken from fully reconstructed MC that has been generated with the fit result to data. We observe small biases in the isobar magnitudes and phases. We correct for these biases by shifting the values of the parameters and assign to this procedure a systematic uncertainty, which corresponds to half the correction combined in quadrature with its error. This uncertainty accounts also for correlations between the signal variables, wrongly reconstructed events, and effects due to the limited sample size (“Fit Bias” in Table V).

From MC we estimate that there are six B background events in our data sample. To determine the bias introduced by these events, we add B background events from MC to our data sample, and fit it with the nominal model. We then assign the observed differences in the observables as a systematic uncertainty (“ B -bkg” in Table V). We assign a systematic uncertainty for all fixed PDF parameters by varying them within their uncertainties according to the covariance matrix.

We vary the histogram PDFs, i.e., the SDP PDF for continuum and the NN PDF for signal (“Discr. Vars” in Table V). The m_{ES} dependence of the SDP PDF for continuum was found to be negligible. We account for differences between simulation and data observed in the control sample $B^0 \rightarrow J/\psi K_S^0$ (“MC-Data” in Table V). These differences were estimated by propagating the dif-




FIG. 6: (color online). One-dimensional scans of $-2\Delta \ln \mathcal{L}$ as a function of magnitudes (left) and phases (right) of the resonances $f_0(980)$, $f_0(1710)$, $f_2(2010)$, and χ_{c0} (top to bottom). The horizontal dashed lines mark the one- and two-standard deviation levels.

ferences, in the control sample, between background-subtracted data and signal MC, into the fit PDFs.

For the branching fraction measurement, we assign a systematic uncertainty due to the error on the calcula-

tion of $N_{B\bar{B}}$ (“ $N_{B\bar{B}}$ ” in Table V) and to the K_S^0 reconstruction efficiency. We correct the K_S^0 reconstruction efficiency by the difference between the efficiency found in a dedicated K_S^0 data sample and that found in simu-

TABLE III: The interference fractions $FF(k, j)$ among the intermediate decay amplitudes for Solution 1. Note that the diagonal elements are those defined in Eq. (28) and detailed in Table II. The lower diagonal elements are omitted since the matrix is symmetric.

	$f_0(980)K_S^0$	$f_0(1710)K_S^0$	$f_2(2010)K_S^0$	NR	$\chi_{c0}K_S^0$
$f_0(980)K_S^0$	0.44	0.07	-0.02	-0.80	0.01
$f_0(1710)K_S^0$		0.07	-0.01	-0.17	-0.0003
$f_2(2010)K_S^0$			0.09	0.02	0.0002
NR				2.16	-0.02
$\chi_{c0}K_S^0$					0.07

TABLE IV: Summary of measurements of branching fractions (\mathcal{B}). The quoted numbers are obtained by multiplying the corresponding fit fraction from Solution 1 by the measured inclusive $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ branching fraction. The first uncertainty is statistical, the second is systematic, and the third represents the signal DP-model dependence.

Mode	$\mathcal{B} [\times 10^{-6}]$
Inclusive $B^0 \rightarrow K_S^0 K_S^0 K_S^0$	$6.19 \pm 0.48 \pm 0.15 \pm 0.12$
$f_0(980)K_S^0, f_0(980) \rightarrow K_S^0 K_S^0$	$2.7^{+1.3}_{-1.2} \pm 0.4 \pm 1.2$
$f_0(1710)K_S^0, f_0(1710) \rightarrow K_S^0 K_S^0$	$0.50^{+0.46}_{-0.24} \pm 0.04 \pm 0.10$
$f_2(2010)K_S^0, f_2(2010) \rightarrow K_S^0 K_S^0$	$0.54^{+0.21}_{-0.20} \pm 0.03 \pm 0.52$
NR, $K_S^0 K_S^0 K_S^0$	$13.3^{+2.2}_{-2.3} \pm 0.6 \pm 2.1$
$\chi_{c0}K_S^0, \chi_{c0} \rightarrow K_S^0 K_S^0$	$0.46^{+0.25}_{-0.17} \pm 0.02 \pm 0.21$

lation. We assign the uncertainty on the correction as a systematic error (“ K_S^0 reco” in Table V).

IV. TIME-DEPENDENT ANALYSIS

In Sec. IV A we describe the proper-time distribution used to extract the time-dependent CP asymmetries. In Sec. IV B we explain the selection requirements used to obtain the signal candidates and suppress backgrounds. In Sec. IV C we describe the fit method and the approach used to account for experimental effects. In Sec. IV D we present the results of the fit and finally, in Sec. IV E we discuss systematic uncertainties in the results.

A. Proper-time distribution

The time-dependent CP asymmetries are functions of the proper-time difference $\Delta t = t_{CP} - t_{\text{tag}}$ between a fully reconstructed $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decay (B_{CP}) and the other B meson decay in the event (B_{tag}), which is partially reconstructed. The observed decay rate is the physical decay rate modified to include tagging imperfections, namely $\langle D \rangle_c$ and ΔD_c ; the former is the rate of correctly assigning the flavor of the B meson, averaged over B^0 and \bar{B}^0 , and the latter is the difference between D_c for B^0 and \bar{B}^0 . The index c denotes different quality

categories of the tag-flavor assignment. Furthermore the decay rate is convolved with the per-event Δt resolution $\mathcal{R}_{\text{sig}}(\Delta t, \sigma_{\Delta t})$, which is described by the sum of three Gaussians and depends on Δt and its error $\sigma_{\Delta t}$. For an event i with tag flavor q_{tag} , one has

$$\begin{aligned} \mathcal{P}_{\text{sig}}^i(\Delta t, \sigma_{\Delta t}; q_{\text{tag}}, c) = & \quad (33) \\ & \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left\{ 1 + q_{\text{tag}} \frac{\Delta D_c}{2} \right. \\ & \left. + q_{\text{tag}} \langle D \rangle_c \left[\mathcal{S} \sin(\Delta m_d \Delta t) - \mathcal{C} \cos(\Delta m_d \Delta t) \right] \right\} \\ & \otimes \mathcal{R}_{\text{sig}}(\Delta t, \sigma_{\Delta t}), \end{aligned}$$

where q_{tag} is defined to be +1 (−1) for $B_{\text{tag}} = B^0$ ($B_{\text{tag}} = \bar{B}^0$), τ_{B^0} is the mean B^0 lifetime, and Δm_d is the mixing frequency [16]. The widths of the B^0 and the \bar{B}^0 are assumed to be the same.

B. Event selection and backgrounds

We reconstruct $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ candidates either from three $K_S^0 \rightarrow \pi^+ \pi^-$ candidates, or from two $K_S^0 \rightarrow \pi^+ \pi^-$ and one $K_S^0 \rightarrow \pi^0 \pi^0$, where the π^0 candidates are formed from pairs of photons. The vertex fit requirements are the same as in the amplitude analysis, and also the requirement that the charged pions of at least one of the K_S^0 have hits in the two inner layers of the vertex tracker. The K_S^0 candidates in the $B^0 \rightarrow 3K_S^0(\pi^+ \pi^-)$ submode must have mass within 12 MeV/ c^2 of the nominal K^0 mass [16] and decay length with respect to the B vertex between 0.2 and 40 cm. In addition, combinatorial background is suppressed in both submodes by imposing that the angle between the momentum vector of each $K_S^0(\pi^+ \pi^-)$ candidate and the vector connecting the beamspot and the $K_S^0(\pi^+ \pi^-)$ vertex is smaller than 0.2 radians. Each K_S^0 decaying to charged pions in the $B^0 \rightarrow 2K_S^0(\pi^+ \pi^-)K_S^0(\pi^0 \pi^0)$ submode is required to have decay length between 0.15 and 60 cm and $\pi^+ \pi^-$ invariant mass less than 11 MeV from the world average K_S^0 mass [16]. The K_S^0 decaying to neutral pions in the $B^0 \rightarrow 2K_S^0(\pi^+ \pi^-)K_S^0(\pi^0 \pi^0)$ submode must have $\pi^0 \pi^0$ invariant mass between 0.48 and 0.52 GeV/ c^2 . Additionally, the neutral pions are selected if they have $\gamma\gamma$ invariant mass between 0.100 and 0.141 GeV/ c^2 and if the photons have energies greater than 50 MeV in the

TABLE V: Summary of systematic uncertainties. The model uncertainty is dominated by the variation of the line shapes due to the contribution of the poorly measured $f_2(2010)$.

Parameter	Fit bias	B -bkg	Discr. Vars	MC-Data	$N_{B\bar{B}}$	K_S^0 reco	Sum	Model
$\mathcal{B}(B^0 \rightarrow K_S^0 K_S^0 K_S^0) [10^{-6}]$	0.011	0.030	0.053	0.015	0.067	0.111	0.145	0.120
FF $f_0(980)$	0.013	0.056	0.006	0.001	-	-	0.058	0.190
FF $f_0(1710)$	0.007	0.001	0.001	0.001	-	-	0.007	0.016
FF $f_2(2010)$	0.005	0.001	0.003	0.001	-	-	0.006	0.084
FF NR	0.024	0.083	0.023	0.001	-	-	0.090	0.344
FF χ_{c0}	0.002	0.000	0.001	0.000	-	-	0.002	0.034
Ph [rad] $f_0(980)$	0.008	0.018	0.014	0.000	-	-	0.024	0.177
Ph [rad] $f_0(1710)$	0.011	0.020	0.001	0.003	-	-	0.023	0.185
Ph [rad] $f_2(2010)$	0.044	0.014	0.004	0.002	-	-	0.046	0.684
Ph [rad] χ_{c0}	0.039	0.011	0.010	0.007	-	-	0.042	0.498

laboratory frame and a lateral energy deposition profile in the electromagnetic calorimeter consistent with that expected for an electromagnetic shower (lateral moment [24] less than 0.55). The fact that we do not model any PDF using sideband data allows a loose requirement on m_{ES} and ΔE in the time-dependent analysis, namely, $5.22 < m_{\text{ES}} < 5.29 \text{ GeV}/c^2$ and $-0.18 < \Delta E < 0.12 \text{ GeV}$. In case of multiple candidates passing the selection, we proceed in the same way as in the amplitude analysis. We use the same NN as in the amplitude analysis to suppress continuum background.

With the above selection criteria, we obtain signal reconstruction efficiencies of 6.7% and 3.1% for the $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ and $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submodes, respectively. These efficiencies are determined from a DP-model MC sample generated using the results of the amplitude analysis. We estimate from MC that 2.1% of the selected signal events are misreconstructed for $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$, while the figure is 2.4% in $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$, and we do not treat these events differently from correctly reconstructed events. Because of the looser requirements, there are more background events from B decays than in the amplitude analysis, in particular in the $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submode. These backgrounds are included in the fit model and are summarized in Table VI. As the analysis is phase-space integrated, we cannot model the χ_{c0} resonance separately, and its contribution to the CP asymmetries could cloud deviations in the charmless contributions. We therefore apply a veto around the invariant mass of this charmonium state.

C. The maximum-likelihood fit

We perform an unbinned extended maximum-likelihood fit to extract the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ event yields along with the \mathcal{S} and \mathcal{C} parameters of the time-dependent analysis.

The fit uses as variables m_{ES} , ΔE , the NN output, Δt , and $\sigma_{\Delta t}$. The selected on-resonance data sample is assumed to consist of signal, continuum background, and backgrounds from B decays. Wrongly reconstructed sig-

nal events are not considered separately. The likelihood function \mathcal{L}_i for event i is the sum

$$\mathcal{L}_i = \sum_j N_j \mathcal{P}_j^i(m_{\text{ES}}, \Delta E, \Delta t, \sigma_{\Delta t}, \text{NN}; q_{\text{tag}}, c, p), \quad (34)$$

where j stands for the species (signal, continuum background, one for each B background category) and N_j is the corresponding yield; q_{tag} , c , and p are the tag flavor, the tagging category, and the physics category, respectively.

To determine q_{tag} and c we use the B flavor-tagging algorithm of Ref. [25]. This algorithm combines several different signatures, such as charges, momenta, and decay angles of charged particles in the event to achieve optimal separation between the two B flavors. This produces six mutually exclusive tagging categories. We also retain untagged events in a seventh category; although these events do not contribute to the measurement of the time-dependent CP asymmetry they do provide additional sensitivity for the measurement of direct CP violation [26].

The two physics categories correspond to $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ and $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ decays. The PDF for species j evaluated for event i is given by the product of individual PDFs:

$$\begin{aligned} \mathcal{P}_j^i(m_{\text{ES}}, \Delta E, \Delta t, \sigma_{\Delta t}, \text{NN}; q_{\text{tag}}, c, p) = \\ \mathcal{P}_j^i(m_{\text{ES}}; p) \mathcal{P}_j^i(\Delta E; p) \mathcal{P}_j^i(\text{NN}; c, p) \mathcal{P}_j^i(\Delta t, \sigma_{\Delta t}; q_{\text{tag}}, c, p). \end{aligned} \quad (35)$$

To take into account the different reconstruction of the two submodes, we use separate PDFs for the two physics categories. Separate NN and Δt PDFs are included for each tagging category within each physics category. The separate Δt PDFs for the two physics categories allow us to fit the \mathcal{S} and \mathcal{C} parameters either separately for the two submodes, or together. The total likelihood is given by

$$\mathcal{L} = \exp\left(-\sum_j N_j\right) \prod_i \mathcal{L}_i. \quad (36)$$

TABLE VI: Summary of B background modes included in the fit model of the time-dependent analysis. The expected number of events takes into account the branching fractions (\mathcal{B}) and efficiencies. In case there is no measurement, the branching fraction of an isospin-related channel is used. All the fixed yields are varied by $\pm 100\%$ for systematic uncertainties.

Submode	Background mode	Varied	$\mathcal{B} [\times 10^{-6}]$	Number of events
$B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$	$K_S^0 K_S^0 K_L^0$	no	2.4	0.71
	$K_S^0 K_S^0 K^{*0}$	no	27.5	9.55
	$K_S^0 K_S^0 K^+$	no	11.5	4.27
	$B^0 \rightarrow \{\text{neutral generic decays}\}$	yes	not applicable	21.7
	$B^+ \rightarrow \{\text{charged generic decays}\}$	yes	not applicable	15.5
$B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$	$K_S^0 K_S^0 K_L^0$	no	2.4	0.67
	$K_S^0 K_S^0 K^{*0}$	no	27.5	5.3
	$K_S^0 K_L^0 K^{*0}$	no	27.5	0.3
	$K_S^0 K_S^0 K^+$	no	11.5	2.9
	$K_S^0 K_S^0 K^{*+}$	no	27.5	7.2
	$B^0 \rightarrow \{\text{neutral generic decays}\}$	yes	not applicable	73.6
	$B^+ \rightarrow \{\text{charged generic decays}\}$	yes	not applicable	73.8

1. Δt PDFs

The signal PDF for Δt is given in Eq. (33). Parameters that depend solely on the tag side of the events (namely, $\langle D \rangle_c$ and ΔD_c) are taken from the analysis of $B \rightarrow c\bar{c}K^{(*)}$ decays [27]. On the other hand, parameters that depend on the signal-side reconstruction, due to the absence of direct tracks from the B decay, cannot be taken from modes that include such direct tracks. This is the case for the parameters that describe the resolution function, which are found in a fit to simulated events. A systematic uncertainty for data-MC differences is assigned using the control sample $B^0 \rightarrow J/\psi K_S^0$, as explained in Sec. IV E.

For continuum events we use a zero-lifetime component. This parametrization is convolved with the same resolution function as for signal, with different parameters that are varied in the fit to data. The parameters of this PDF are not separated in the tagging categories. The small contribution from $e^+e^- \rightarrow c\bar{c}$ events is well described by the tails of the resolution function. For B background events we use the signal PDF, with resolution parameters from the BABAR $B \rightarrow c\bar{c}K^{(*)}$ analysis [27]. The parameters \mathcal{S} and \mathcal{C} are set to zero and varied to assign a systematic uncertainty.

2. Description of the other variables

The m_{ES} and ΔE distributions of signal events are parametrized by an asymmetric Gaussian with power-law tails, as given in Eq. (24), and, for m_{ES} , a small additional component, parameterized by an ARGUS shape function [20], to correctly describe misreconstructed events. The means in these two PDFs for $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ events are allowed to vary in the fit to data, and the other parameters are taken from MC simulation. For the NN distributions of signal we use his-

togram PDFs taken from MC simulation for each physics and tagging category. The m_{ES} , ΔE , and NN PDFs for continuum events are parametrized by an ARGUS shape function, a straight line and the sum of power functions from Eq. (25), respectively. All continuum parameters, except for c_2 and c_3 of the NN PDF, are allowed to vary in the fit. All the fixed parameters are varied, within the uncertainties found in a fit to sideband data, to estimate systematic errors.

All the B background PDFs are described by fixed histograms taken from MC simulation.

D. Results

The maximum-likelihood fit of 3261 candidates in the $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ submode and 7209 candidates in the $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submode results in the event yields detailed in Table VII.

TABLE VII: Event yields for the different event species, resulting from the maximum-likelihood fit for the time-dependent analysis. “ B^+B^- ($B^0\bar{B}^0$) bkg” represents background from charged (neutral) B decays. Quoted uncertainties are statistical only.

Species	$3K_S^0(\pi^+\pi^-)$	$2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$
Signal	201^{+16}_{-15}	62^{+13}_{-12}
Continuum	3086^{+56}_{-54}	7086^{+85}_{-83}
B^+B^- bkg	-54^{+29}_{-24}	45^{+34}_{-30}
$B^0\bar{B}^0$ bkg	9^{+31}_{-30}	4^{+38}_{-29}

The fit result for the time-dependent CP -violation parameters \mathcal{S} and \mathcal{C} is

$$\begin{aligned}\mathcal{S} &= -0.94^{+0.24}_{-0.21}, \\ \mathcal{C} &= -0.17 \pm 0.18,\end{aligned}$$

where the uncertainties are statistical only. The correlation between \mathcal{S} and \mathcal{C} is -0.16 . We use the fit result to create *sPlots* of the signal distributions of Δt , the time-dependent asymmetry, and the discriminating variables. Figure 7 shows the Δt *sPlots* for the combined fit result and for the individual submodes. Figure 8 shows the signal distributions and Fig. 9 the continuum background distributions of the discriminating variables. The distributions shown in these three figures illustrate the good agreement between the data and the fit model.

We scan the statistical-only likelihood of the \mathcal{S} parameter for both submodes and for the combined fit. The result, on the left-hand side of Fig. 10, shows a sizable difference between the \mathcal{S} values for the two submodes; the level of consistency, conservatively estimated from the sum of the two individual likelihood scans, is approximately 2.6σ (a p -value of 1.0% with 2 degrees of freedom). This value is obtained including only the dominant statistical uncertainty and neglecting the small correlation between the CP -violation parameters. The results obtained when \mathcal{S} and \mathcal{C} are allowed to vary individually for each of the submodes are $\mathcal{S} = -1.42^{+0.27}_{-0.24}$, $\mathcal{C} = -0.14^{+0.17}_{-0.17}$ for $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ and $\mathcal{S} = 0.40^{+0.56}_{-0.57}$, $\mathcal{C} = 0.19^{+0.42}_{-0.43}$ for $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$. In both cases the quoted uncertainties are statistical only.

As there is some correlation between the \mathcal{S} and \mathcal{C} parameters, we perform a two-dimensional statistical likelihood scan of the combined likelihood, which is then convolved by the systematic uncertainties on \mathcal{S} and \mathcal{C} (systematic uncertainties are discussed in Sec. IV E.) The result is shown on the right-hand side of Fig. 10. We find that CP conservation is excluded at 3.8 standard deviations, and thus, for the first time, we measure an evidence of CP violation in $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays. The difference between our result and that from $B^0 \rightarrow c\bar{c}K^{(*)}$ is less than 2 standard deviations. The scan also shows that the result is close to the physical boundary, given by the constraint $\mathcal{S}^2 + \mathcal{C}^2 \leq 1$.

E. Systematic uncertainties

The systematic uncertainties are summarized in Table VIII. The “MC_{stat}” uncertainty accounts for the limited size of the simulated data samples used to create the PDFs. The “ B_{reco} ” uncertainty propagates the experimental uncertainty in the measurement of tag-side-related quantities taken from [27] to our measurement. The “ B -bkg” contribution results from the uncertainty in the CP content and the branching fractions of fixed yields in the model of background from B decays. The dominant “MC-Data: Δt ” systematic uncertainty is due to possible differences between data and simulation concerning the procedure used to obtain the signal B decay vertex from tracks originating from K_S^0 decays. We quantify this uncertainty using the control sample $B^0 \rightarrow J/\psi K_S^0$ by comparing the difference between Δt values obtained with and without the J/ψ in

data and simulation. We then propagate the observed differences and their uncertainties to the resolution function. We use this new resolution function to refit the data and obtain an estimate of the effect on \mathcal{S} and \mathcal{C} . We also use the samples $B^0 \rightarrow J/\psi K_S^0(\pi^+\pi^-)$ and $B^0 \rightarrow J/\psi K_S^0(\pi^0\pi^0)$ to estimate simulation-data differences for the other variables in the submodes $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ and $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$, respectively. This contribution is referred to as “MC-Data: Discr. Vars” in Table VIII. The “Fit Bias” uncertainty is evaluated using fits to fully reconstructed MC samples. It accounts for effects from wrongly reconstructed events and correlations between fit variables. The “Veto” uncertainty is related to the veto on the invariant mass. It is evaluated using events that pass the veto in pseudo-experiments studies. Finally, the “Misc” uncertainty includes contributions from doubly-Cabibbo-suppressed decays, silicon vertex tracker alignment, and the uncertainties in the boost of the $\Upsilon(4S)$. These contributions are taken from the BABAR $B \rightarrow c\bar{c}K^{(*)}$ analysis [27].

V. SUMMARY

We have performed the first amplitude analysis of $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ events and measured the total inclusive branching fraction to be $(6.19 \pm 0.48 \pm 0.15 \pm 0.12) \times 10^{-6}$, where the first uncertainty is statistical, the second is systematic, and the third represents the signal DP-model dependence. We have identified the dominant contributions to the DP to be from $f_0(980)$, $f_0(1710)$, $f_2(2010)$, and a nonresonant component, and measured the individual fit fractions and phases of each component. We do not observe any significant contribution from the so-called $f_X(1500)$ resonance seen in, for example, $B^+ \rightarrow K^+ K^- K^+$ [6]. The peak in the invariant mass between 1.5 and 1.6 GeV/ c^2 can be described by the interference between the $f_0(1710)$ resonance and the nonresonant component. We see some hints from the $f_2'(1525)$ and $f_0(1500)$ resonances that could also contribute to this structure, but due to limited sample size we cannot make a significant statement. Future investigations of the KK system could shed more light on the situation. Furthermore we have performed an update of the phase-space-integrated time-dependent analysis of the same decay mode, using $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ and $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ decays, with the final BABAR dataset. We measure the CP -violation parameters to be $\mathcal{S} = -0.94^{+0.24}_{-0.21} \pm 0.06$ and $\mathcal{C} = -0.17 \pm 0.18 \pm 0.04$, where the first quoted uncertainty is statistical and the second is systematic. These measured values are consistent with and supersede those reported in Ref. [3]. They are compatible within two standard deviations with those measured in tree-dominated modes such as $B^0 \rightarrow J/\psi K_S^0$, as expected in the SM. For the first time, we report evidence of CP violation in $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decays; CP conservation is excluded at 3.8 standard deviations including systematic uncertainties.




FIG. 7: (color online). Signal $sPlots$ (points with error bars) and PDFs (histograms) of Δt (left) and the derived asymmetry (right) for the $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ submode (top), the $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submode (middle), and the combined fit (bottom). In the Δt distributions on the left-hand side, points marked with \times and solid lines correspond to decays where B_{tag} is a B^0 meson; points marked with \circ and dashed lines correspond to decays where B_{tag} is a \bar{B}^0 meson. Points of the asymmetry $sPlots$ that are outside the range of a figure are marked by arrows.

TABLE VIII: Summary of systematic uncertainties on the \mathcal{S} and \mathcal{C} parameters.

Source	\mathcal{S}	\mathcal{C}
MC _{stat}	0.002	0.001
B_{reco}	0.004	0.003
B -bkg	0.032	0.012
MC-Data: Δt	0.045	0.027
MC-Data: Discr. Vars	0.021	0.004
Fit Bias	0.022	0.018
Veto	0.006	0.004
Misc	0.004	0.015
Sum	0.064	0.038




FIG. 8: (color online). Signal s Plots (points with error bars) and PDFs (histograms) of the discriminating variables: m_{ES} (top), ΔE (middle), and the NN output (bottom) for the $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ submode (left) and for the $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submode (right). Below each bin are shown the residuals, normalized in error units. The horizontal dotted and full lines mark the one- and two-standard deviation levels, respectively.




FIG. 9: (color online). Continuum s Plots (points with error bars) and PDFs (histograms) of m_{ES} , ΔE , the NN output, and Δt (top to bottom). Plots on the left-hand side correspond to the $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ submode, and on the right-hand side to the $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submode. In the Δt distributions, points marked with \times and solid lines correspond to decays where B_{tag} is a B^0 meson; points marked with \circ and dashed lines correspond to decays where B_{tag} is a \bar{B}^0 meson. Below each bin are shown the residuals, normalized in error units. The horizontal dotted and full lines mark the one- and two-standard deviation levels, respectively.

ACKNOWLEDGMENTS

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminos-

ity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions




FIG. 10: (color online). One-dimensional statistical scan of $-2\Delta\ln\mathcal{L}$ as a function of \mathcal{S} (left) and the two-dimensional scan, including systematic uncertainty, as a function of \mathcal{S} and \mathcal{C} (right). In the left-hand plot, red points marked with \times correspond to the $B^0 \rightarrow 3K_S^0(\pi^+\pi^-)$ submode, blue points marked with \circ to the $B^0 \rightarrow 2K_S^0(\pi^+\pi^-)K_S^0(\pi^0\pi^0)$ submode, and black points marked with $*$ to the combined fit. In the right-hand plot, the gray scale is given in units of $\sqrt{-2\Delta\ln\mathcal{L}}$. The result of the *BABAR* analyses of $B^0 \rightarrow c\bar{c}K^{(*)}$ decays [27] is indicated as a white ellipse and the physical boundary ($\mathcal{S}^2 + \mathcal{C}^2 \leq 1$) is marked as a gray line. The scan appears to be trimmed on the lower left since the PDF becomes negative outside the physical region (i.e., the white region does not indicate that the scan flattens out at 5σ).

wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), the Commissariat à l'Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nu-

cleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Education and Science of the Russian Federation, Ministerio de Ciencia e Innovación (Spain), and the Science and Technology Facilities Council (United Kingdom). Individuals have received support from the Marie-Curie IEF program (European Union), the A. P. Sloan Foundation (USA) and the Binational Science Foundation (USA-Israel).

-
- [1] Y. Grossman and M. P. Worah, *Phys. Lett. B* **395**, 241 (1997).
 - [2] H.-Y. Cheng, C.-K. Chua, and A. Soni, *Phys. Rev. D* **72**, 094003 (2005).
 - [3] B. Aubert et al. (*BABAR*), *Phys. Rev. D* **76**, 091101 (2007).
 - [4] K.-F. Chen et al. (*Belle*), *Phys. Rev. Lett.* **98**, 031802 (2007).
 - [5] T. Gershon and M. Hazumi, *Phys. Lett. B* **596**, 163 (2004).
 - [6] B. Aubert et al. (*BABAR*), *Phys. Rev. D* **74**, 032003 (2006).
 - [7] B. Aubert et al. (*BABAR*), *Phys. Rev. Lett.* **99**, 161802 (2007).
 - [8] B. Aubert et al. (*BABAR*) (2008), presented at ICHEP 2008, arXiv:0808.0700.
 - [9] A. Garmash et al. (*Belle*), *Phys. Rev. D* **71**, 092003 (2005).
 - [10] Y. Nakahama et al. (*Belle*), *Phys. Rev. D* **82**, 073011 (2010).
 - [11] B. Aubert et al. (*BABAR*), *Phys. Rev. Lett.* **99**, 221801 (2007).
 - [12] B. Aubert et al. (*BABAR*), *Phys. Rev. D* **79**, 051101 (2009).
 - [13] N. Rey-Le Lorier and D. London, *Phys. Rev. D* **85**, 016010 (2012).
 - [14] B. Aubert et al. (*BABAR*), *Nucl. Instrum. Methods Phys. Res. A* **479**, 1 (2002).
 - [15] J. Blatt and V. E. Weisskopf, *Theoretical Nuclear Physics* (J. Wiley, New York, 1952).
 - [16] C. Amsler et al. (Particle Data Group), *Phys. Lett. B* **667**, 1 (2008).
 - [17] S. M. Flatte, *Phys. Lett. B* **63**, 224 (1976).
 - [18] M. Ablikim et al. (*BES*), *Phys. Lett. B* **607**, 243 (2005).
 - [19] P. Gay, B. Michel, J. Proriot, and O. Deschamps (1995), prepared for 4th International Workshop on Software Engineering and Artificial Intelligence for High-energy and Nuclear Physics (AIHENP 95), Pisa, Italy, 1995.
 - [20] H. Albrecht et al. (*ARGUS*), *Z. Phys. C* **48**, 543 (1990).
 - [21] M. Pivk and F. R. Le Diberder, *Nucl. Instrum. Methods Phys. Res. A* **555**, 356 (2005).
 - [22] B. Aubert et al. (*BABAR*), *Phys. Rev. D* **80**, 112001 (2009).
 - [23] P. del Amo Sanchez et al. (*BABAR*), *Phys. Rev. D* **83**, 052001 (2011).
 - [24] A. Drescher et al., *Nucl. Instrum. Methods Phys. Res. A* **237**, 464 (1985).
 - [25] B. Aubert et al. (*BABAR*), *Phys. Rev. Lett.* **94**, 161803 (2005).
 - [26] S. Gardner and J. Tandean, *Phys. Rev. D* **69**, 034011 (2004).
 - [27] B. Aubert et al. (*BABAR*), *Phys. Rev. D* **79**, 072009 (2009).

(2009).